

CMB constraint on non-Gaussianity in isocurvature perturbations

Chiaki Hikage^a, Masahiro Kawasaki^{b,c}, Toyokazu Sekiguchi^d and Tomo Takahashi^e

^a*Kobayashi-Maskawa Institute, Nagoya University, Nagoya 464-8602, Japan*

^b*Institute for Cosmic Ray Research, University of Tokyo, Kashiwa 277-8582, Japan*

^c*Kavli Institute for the Physics and Mathematics of the Universe, University of Tokyo, Kashiwa 277-8568, Japan*

^d*Department of Physics and Astrophysics, Nagoya University, Nagoya 464-8602, Japan*

^e*Department of Physics, Saga University, Saga 840-8502, Japan*

Abstract

We study the CMB constraint on non-Gaussianity in CDM isocurvature perturbations. Non-Gaussian isocurvature perturbations can be produced in various models at the very early stage of the Universe. Since the isocurvature perturbations little affect the structure formation at late times, CMB is the best probe of isocurvature non-Gaussianity at least in the near future. In this paper, we focus on uncorrelated isocurvature perturbations and constrain their non-Gaussianity. For this purpose, we employ several state-of-art techniques for the analysis of CMB data and simulation. We use the WMAP 7 year data of temperature anisotropy. When the adiabatic perturbations are assumed to be Gaussian, we obtained a constraint on the isocurvature non-Gaussianity $\alpha^2 f_{\text{NL}}^{(\text{ISO})} = 40 \pm 66$ for the scale invariant isocurvature power spectrum, where α is the ratio of the power spectrum of isocurvature perturbations to that of the adiabatic ones. When we assume that the adiabatic perturbations can also be non-Gaussian, we obtain $f_{\text{NL}} = 38 \pm 24$ and $\alpha^2 f_{\text{NL}}^{(\text{ISO})} = -8 \pm 72$. We also discuss implications our results on the axion CDM isocurvature model.

1 Introduction

The adiabaticity, or isocurvature mode of primordial density fluctuations is one of important probes of cosmology in various respects. Although current cosmological observations suggest that primordial density fluctuations are almost adiabatic and they give severe constraints on the size of isocurvature fluctuations, some fraction of their contribution is still allowed [1]. Residual isocurvature fluctuations can be generated when there exist multiple components with different origins, which are associated with dark matter, baryon and neutrino [2]^{#1}. Such examples include axion [5, 6, 7, 8, 9, 10, 11, 12] and Affleck-Dine baryogenesis [13, 14, 15], where cold dark matter (CDM) and baryon isocurvature modes can be respectively generated. These isocurvature modes are basically uncorrelated with adiabatic ones, however, when one considers a scenario where a light scalar field other than the inflaton is responsible for (adiabatic) density fluctuation such as the curvaton model [16, 17, 18], isocurvature perturbations can be correlated with the adiabatic ones and be easily generated, depending on how and when CDM and baryon are created [19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31]. In addition to above mentioned examples, a variety of models with isocurvature fluctuations has been extensively studied in the literature, hence the information on isocurvature fluctuations would give a lot of insight on various aspects of cosmology, particularly on models of dark matter and baryogenesis as well as those of the early Universe.

Although the theoretical works on non-Gaussianity from isocurvature fluctuations have been relatively well studied [32, 33, 34, 35, 36, 37], observational constraints on them have not been explored much^{#2}. In particular, we cannot find any work investigating this issue by using the actual data except Ref. [48], where the constraint has been studied with Minkowski functionals using the WMAP 3 data, although there are a few papers which elaborate its future CMB constraints [49, 50, 51]. In the light that we now have precise cosmological data to explore non-Gaussianity as seen from the counterpart for adiabatic ones, non-Gaussianity in isocurvature fluctuations should be pursued more.

In this paper, we investigate the optimal constraint on non-Gaussianity in CDM (baryon) isocurvature fluctuations from CMB bispectrum estimator. Although, as mentioned above, isocurvature fluctuations can be correlated with adiabatic ones in some cases and there are also other kinds of modes such as neutrino one, we in this paper present the methodology of our analysis and concentrate to report the constraint on the CDM (baryon) uncorrelated type. Constraints on other types such as correlated ones and neutrino modes will be reported in a forthcoming paper [52].

The organization of this paper is as follows. In the next section, we give the formalism to discuss non-Gaussianity in models with isocurvature fluctuations and also set our notation. In Section 3, we describe our analysis method to obtain the constraint on non-Gaussianity from isocurvature fluctuations. Then in Section 4, we present our results

^{#1} Isocurvature fluctuations in dark radiation have also been studied in Refs. [3, 4].

^{#2} At linear order, there are a lot of works which study the observational constraints [38, 39, 40, 41, 42, 43, 44, 45, 46, 47].

on the constraint. As an application of our results, we consider constraints on the axion isocurvature model in Section 5. The final section is devoted to the conclusion of this paper.

2 Model of non-Gaussian perturbations and CMB signature

We consider primordial curvature and CDM isocurvature perturbations, respectively denoted as ζ and S , in the following form:

$$\zeta(\vec{x}) = \zeta_G(\vec{x}) + \frac{3}{5} f_{\text{NL}} (\zeta_G(\vec{x})^2 - \langle \zeta_G(\vec{x})^2 \rangle), \quad (1)$$

$$S(\vec{x}) = S_G(\vec{x}) + f_{\text{NL}}^{(\text{ISO})} (S_G(\vec{x})^2 - \langle S_G(\vec{x})^2 \rangle), \quad (2)$$

where ζ_G and S_G are Gaussian parts of the primordial curvature and isocurvature perturbations, respectively. f_{NL} and $f_{\text{NL}}^{(\text{ISO})}$ are the non-linearity parameters of the curvature and isocurvature perturbations, respectively. In the following, we denote these primordial perturbations with $X^A(\vec{x})$. Then Eqs. (1) and (2) can be recast into

$$X^A(\vec{x}) = X_G^A(\vec{x}) + f_{\text{NL}}^A (X_G^A(\vec{x})^2 - \langle X_G^A(\vec{x})^2 \rangle), \quad (3)$$

where X_G^A is the Gaussian part of X^A . Here we defined a non-linearity parameter f_{NL}^A , which is related to the adiabatic and isocurvature ones via $f_{\text{NL}}^\zeta = \frac{3}{5} f_{\text{NL}}$ and $f_{\text{NL}}^S = f_{\text{NL}}^{(\text{ISO})}$.

In this paper, we consider uncorrelated isocurvature perturbations, so that $\langle \zeta_G S_G \rangle = 0$. Thus only correlation functions which contain either ζ or S are non-zero. In terms of the Fourier components of X^A , the bispectrum from either the primordial curvature or isocurvature perturbations is

$$\langle X^A(\vec{k}_1) X^A(\vec{k}_2) X^A(\vec{k}_3) \rangle = 2 f_{\text{NL}}^A (P_{X_G^A}(k_1) P_{X_G^A}(k_2) + (2 \text{ perms})), \quad (4)$$

where $P_{X_G^A}(k)$ is the power spectrum of the Gaussian perturbations X_G^A . Here we neglected loop contributions although we take those into account in Sec. 5.

Neglecting the secondary non-Gaussianities arising from the second or higher order cosmological perturbation theory, the harmonic coefficients of the CMB temperature anisotropy from primordial perturbations X^A can be given as

$$a_{lm}^A = 4\pi(-i)^l \int \frac{dk^3}{(2\pi)^3} g_l^A(k) X^A(\vec{k}) Y_{lm}(\hat{k}), \quad (5)$$

where $g_l^A(k)$ is the temperature transfer function for the primordial perturbations X^A . The total CMB anisotropy is the sum of those from the curvature and isocurvature perturbations, i.e. $a_{lm} = a_{lm}^\zeta + a_{lm}^S$.

Since ζ and S are uncorrelated we need to consider the polyspectra which contain only either ζ or S . Then the total polyspectra are the sum of those from each perturbation. Angular power spectrum C_l^A , which is defined by $\langle a_{lm}^A a_{l'm'}^{A*} \rangle = C_l^A \delta_{ll'} \delta_{mm'}$ can be given as

$$C_l^A = \frac{2}{\pi} \int dk k^2 g_l^A(k)^2 P_{X_G^A}(k). \quad (6)$$

The reduced bispectrum $b_{l_1 l_2 l_3}$ is defined by

$$\langle a_{l_1 m_1} a_{l_2 m_2} a_{l_3 m_3} \rangle = b_{l_1 l_2 l_3} \mathcal{G}_{m_1 m_2 m_3}^{l_1 l_2 l_3}, \quad (7)$$

where $\mathcal{G}_{m_1 m_2 m_3}^{l_1 l_2 l_3}$ is the Gaunt integral, which can be written in terms of the Wigner-3j symbol as

$$\mathcal{G}_{m_1 m_2 m_3}^{l_1 l_2 l_3} = \sqrt{\frac{(2l_1 + 1)(2l_2 + 1)(2l_3 + 1)}{4\pi}} \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \begin{pmatrix} l_1 & l_2 & l_3 \\ 0 & 0 & 0 \end{pmatrix}. \quad (8)$$

Given a bispectrum of primordial perturbations in Eq. (4), reduced CMB bispectrum from each of the primordial perturbations can be given as

$$b_{l_1 l_2 l_3}^A = 2f_{\text{NL}}^A \int dr r^2 (\alpha_{l_1}^A(r) \beta_{l_2}^A(r) \beta_{l_3}^A(r) + (2 \text{ perms})), \quad (9)$$

where $\alpha_l^A(r)$ and $\beta_l^A(r)$ are

$$\alpha_l^A(r) = \frac{2}{\pi} \int dk k^2 g_l^A(k) j_l(kr), \quad (10)$$

$$\beta_l^A(r) = \frac{2}{\pi} \int dk k^2 g_l^A(k) j_l(kr) P_{X_G^A}(k). \quad (11)$$

For later convenience, we introduce a normalized bispectrum

$$\hat{b}_{l_1 l_2 l_3}^A \equiv b_{l_1 l_2 l_3}^A / f_{\text{NL}}^A. \quad (12)$$

The amplitude of isocurvature power spectrum is constrained from CMB angular power spectrum. In Ref. [1], it is shown that the WMAP observation of CMB gives an upper bound on the fraction of isocurvature power spectrum in the total one

$$\alpha \equiv \frac{P_S}{P_\zeta} < 0.15 \quad (13)$$

at 95 % confidence level.

Note that the bispectrum is proportional to $\alpha^2 f_{\text{NL}}^{(\text{ISO})}$. This can be seen from Eqs. (9) and (11). Hence we report our constraint on non-Gaussianity in isocurvature fluctuations for the combination of $\alpha^2 f_{\text{NL}}^{(\text{ISO})}$.

3 Analysis method

Here we describe our analysis method to give the constraint on the non-linearity parameters from CMB data. Our method basically follows the one in Ref. [54].

3.1 Estimator of non-linearity parameters

In the limit of small non-linearity parameter f_{NL}^A , the effect of the deviation from Gaussianity manifests in the CMB bispectrum, so that the optimal estimator can be constructed from the three point function of CMB anisotropy [55]. We adopt the following cubic estimator of non-linearity parameters

$$\begin{pmatrix} \hat{f}_{\text{NL}}^\zeta \\ \hat{f}_{\text{NL}}^S \end{pmatrix} = \begin{pmatrix} \langle S_{\text{prim}}^\zeta \rangle_{f_{\text{NL}}^\zeta=1} & \langle S_{\text{prim}}^\zeta \rangle_{f_{\text{NL}}^S=1} \\ \langle S_{\text{prim}}^S \rangle_{f_{\text{NL}}^\zeta=1} & \langle S_{\text{prim}}^S \rangle_{f_{\text{NL}}^S=1} \end{pmatrix}^{-1} \begin{pmatrix} S_{\text{prim}}^\zeta \\ S_{\text{prim}}^S \end{pmatrix}, \quad (14)$$

where the square bracket with subscript $f_{\text{NL}}^A = 1$ indicates an ensemble average over simulations with non-zero non-linearity parameter shown in the subscript with the other non-linearity parameter being fixed to zero. S_{prim}^A is the cubic statistics [55, 56, 57, 58], which is given by

$$S_{\text{prim}}^A = \frac{1}{6} \sum_{\{lm\}} \hat{b}_{l_1 l_2 l_3}^A \mathcal{G}_{m_1 m_2 m_3}^{l_1 l_2 l_3} [\tilde{a}_{l_1 m_1} \tilde{a}_{l_2 m_2} \tilde{a}_{l_3 m_3} - 3 \tilde{a}_{l_1 m_1} \langle \tilde{a}_{l_2 m_2} \tilde{a}_{l_3 m_3} \rangle_0], \quad (15)$$

where the square bracket with subscript 0 indicates an ensemble average over Gaussian simulations. $\tilde{a}_{lm}$ is a harmonic coefficient obtained from observed (or simulated) data maps with suitable weighting, which will be discussed in Section 3.3. Eq. (14) can be schematically represented as $\hat{f}_{\text{NL}}^A = \sum_B \langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1}^{-1} S_{\text{prim}}^B$.

In our analysis, to estimate $\langle \tilde{a}_{lm} \tilde{a}_{l'm'} \rangle$ we accumulated 500 Monte Carlo (MC) samples. This number is sufficient, since the uncertainty in estimation of S_{prim}^A due to this limitation of MC samples is much smaller than the Gaussian error $\sqrt{\langle S_{\text{prim}}^A S_{\text{prim}}^A \rangle_0}$.

Assuming the Gaussianity for X^A , we estimate the covariance of the estimator $\hat{f}_{\text{NL}}^A$,

$$\begin{aligned} \langle \hat{f}_{\text{NL}}^A \hat{f}_{\text{NL}}^B \rangle_0 &= \sum_{A'B'} \langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^{A'}=1}^{-1} \langle S_{\text{prim}}^{A'} S_{\text{prim}}^{B'} \rangle_0 \langle S_{\text{prim}}^B \rangle_{f_{\text{NL}}^{B'}=1}^{-1} \\ &= \langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^A=1}^{-1}. \end{aligned} \quad (16)$$

Here we used the relation $\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1} = \langle S_{\text{prim}}^A S_{\text{prim}}^B \rangle_0$. See Appendix A for the derivation.

The estimator in Eq. (14) can be regarded as the generalization of the fast estimator in Ref. [55] to the case where the initial perturbations are mixture of non-Gaussian adiabatic and uncorrelated isocurvature perturbations. When we assume that either of adiabatic

or uncorrelated isocurvature perturbations are non-Gaussian and the other is Gaussian, Eq. (14) can be reduced to

$$\hat{f}_{\text{NL}}^A = S_{\text{prim}}^A / \langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^A=1}, \quad (17)$$

where the subscript A indicates the perturbations which are assumed to be non-Gaussian. In this case, the variance of $\hat{f}_{\text{NL}}^A$ is given by $1 / \langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^A=1}$.

We note that there is a difficulty in computing the normalization factor $\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1}$. If we naively evaluate Eq. (15) with small but non-zero f_{NL}^A and average them, a substantial number of simulations are required due to a large Gaussian fluctuation. Instead, we divide $\tilde{a}_{lm}$ into its Gaussian and non-Gaussian parts and evaluate S_{prim}^A without terms which are to vanish by averaging. The details are presented in Appendix B. This treatment also removes the need for setting a non-zero fiducial α .

3.2 Non-Gaussian CMB simulation

In order to determine the normalization $\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1}$, we need to simulate non-Gaussian CMB maps. With the Fourier transformation, Eq. (5) can be rewritten as

$$a_{lm}^A = \int dr r^2 \alpha_l^A(r) \int d\hat{r} Y_{lm}^*(\hat{r}) X^A(\vec{r}). \quad (18)$$

In the case of local-type non-Gaussianity, Eq. (18) allows us to simulate the non-Gaussian CMB maps exactly [59, 60]. We adopt the fast method developed in Ref. [61], which enables simulations of the local-type non-Gaussianity with sufficient speed. The key technique in the method of Ref. [61] is that the line of sight integral in Eq. (18) is evaluated by the Gaussian quadrature with optimized nodes and weights. In our simulation, we optimized the nodes so that the mean square of the error in simulated maps a_{lm} at each multipole (lm) should be less than 0.01. For $l_{\text{max}} = 1024$, we found that this level of accuracy requires 42 and 15 nodes for curvature and isocurvature perturbations, respectively.

3.3 Optimally-weighted CMB maps

In estimating $\hat{f}_{\text{NL}}^A$, we need to suitably weight observed (and simulated) maps a_{lm} to obtain $\tilde{a}_{lm}$, in order to take into account the sensitivity and resolution of the survey. As discussed in Ref. [54], the optimal weight is the inverse of the variance. This can be schematically represented as $\tilde{a}_{lm} = [C^{-1}a]_{lm}$, where $C = C_S + C_N$ is the covariance matrix, with those of signal and noise being denoted as C_S and C_N , respectively. While C_S is diagonal in the harmonic space, C_N is in general complicated for real observations. In particular, WMAP and many CMB surveys have multiple channels, so that we need to take into account different beam functions, inhomogeneous noises, and a sky cut. In such a case, $\tilde{a}_{lm}$ can be given in an implicit form as

$$(C_S^{-1} + C_N^{-1})C_S\tilde{a} = C_N^{-1}a, \quad (19)$$

where $C_N^{-1} = \sum_i b^{(i)} C_N^{(i)-1} b^{(i)}$ and $C_N^{-1} a = \sum_i C_N^{(i)-1} a^{(i)}$, with the subscript i indicating a channel of the survey. $b^{(i)}$ here is the beam function.

Since the matrix of $(C_S^{-1} + C_N^{-1})$ is dense in both the harmonic and pixel spaces, direct implementation of inversion $(C_S^{-1} + C_N^{-1})^{-1}$ is substantially prohibited. Instead, Eq. (19) can be solved by the conjugate gradient (CG) method with preconditioning [62]. How fast a CG method converges crucially depends on a choice of a pre-conditioner. We adopt a fast method developed in Ref. [63], which makes use of the multi-grid preconditioning. This method also allows to marginalize over the amplitude of components which are proportional to template maps $\{\tau\}$. To do this, C_N^{-1} is replaced with $C_N^{-1} - \sum_{ab} C_N^{-1} \tau_a (\tau_a C_N^{-1} \tau_b)^{-1} C_N^{-1} \tau_b$, where the subscripts a and b indicate template components. For the details of the method, we refer to Refs. [63, 54].

4 Result

In our analysis, we assume a flat power-law Λ CDM model as a fiducial model and adopt the mean values for cosmological parameters from the WMAP 7 year data alone [1]. We also assume that the power spectrum of isocurvature perturbations is in a power-law form $P_{S_G}(k) \propto k^{n_{\text{iso}}-4}$, and with regard to the fiducial value of the spectral index n_{iso} we consider two different cases, $n_{\text{iso}} = 0.963 (= n_{\text{adi}})$ and $n_{\text{iso}} = 1$. The transfer function of CMB is computed using the CAMB code [64]. We combine the foreground-cleaned maps of V and W bands of the WMAP 7 year data [65, 66]^{#3} with a resolution $N_{\text{side}} = 512$ in the HEALPix pixelization scheme [67]^{#4}. We adopt the KQ75y7 mask [66], which cuts 28.4 % of the sky. We also set the maximum multipole l_{max} to 1024 in our analysis. We marginalize the amplitudes of the monopole $l = 0$ and dipoles $l = 1$ as default. We also optionally marginalize the amplitudes of the Galactic foreground components at large angular scales using the synchrotron, free-free and dust emission templates from Ref. [66]. Our results are summarized in Table 1.

To check our analysis method, we have evaluated a constraint on the non-linearity parameter for adiabatic perturbations assuming isocurvature perturbations being absent, and compare it with those in a previous study. In this case, we obtain $f_{\text{NL}} = 31 \pm 21$ at 1σ level without template marginalization of the Galactic foregrounds. Without template marginalization, our central value by about 0.5σ deviates from that of Ref. [1], which gives $f_{\text{NL}} = 42 \pm 21$ from the foreground cleaned V+W band data with a resolution $N_{\text{side}} = 1024$. Since there are substantial differences in our analysis and that of Ref. [1], including different fiducial cosmological models and resolutions of the maps used, we believe that this level of difference is acceptable and our result is consistent with the previous study. With template marginalization, we obtain $f_{\text{NL}} = 32 \pm 21$, which is exactly the same as the

^{#3}<http://lambda.gsfc.nasa.gov>

^{#4}<http://healpix.jpl.nasa.gov>

	setups	f_{NL}	$\alpha^2 f_{\text{NL}}^{(\text{ISO})}$
$n_{\text{iso}} = 0.963$	w/o template marginalization	31 ± 21 (36 ± 23)	5 ± 63 (-39 ± 69)
	w/ template marginalization	32 ± 21 (32 ± 23)	40 ± 63 (0 ± 70)
$n_{\text{iso}} = 1$	w/o template marginalization	31 ± 21 (34 ± 23)	19 ± 62 (-22 ± 70)
	w/ template marginalization	32 ± 21 (38 ± 24)	40 ± 66 (-8 ± 72)

Table 1: Constraints on f_{NL} and $\alpha^2 f_{\text{NL}}^{(\text{ISO})}$ for different setups. We adopted four different setups regarding the fiducial value of n_{iso} and template marginalization of the Galactic foregrounds. Constraints without parenthesis are estimated by fixing the other non-linearity parameter to zero. On the other hand, ones with parenthesis are estimated by marginalizing over the other non-linearity parameter.

one given by the WMAP group [1]. We found that template marginalization little affects constraints on f_{NL} .

Now we present constraints on non-Gaussianity in uncorrelated isocurvature perturbations. We first assume that the adiabatic perturbations are Gaussian and fix f_{NL} to be zero. For the cases of $n_{\text{iso}} = 0.963$ and $n_{\text{iso}} = 1$, we respectively obtain $\alpha^2 f_{\text{NL}}^{(\text{ISO})} = 5 \pm 63$ and 19 ± 62 at 1σ level without template marginalization. With template marginalization, these change to 40 ± 63 and 40 ± 66 . We found that the constraints on $\alpha^2 f_{\text{NL}}^{(\text{ISO})}$ are not strongly affected by the fiducial value of n_{iso} . On the other hand, the constraints are more or less dependent on the treatment of the Galactic foregrounds. The central values can change by 0.5σ while the error remains almost unchanged. However, this shows that the effects of residual foregrounds are not severe.

When we assume that both adiabatic and isocurvature perturbations can be non-Gaussian, we give a joint constraint on $(f_{\text{NL}}, \alpha^2 f_{\text{NL}}^{(\text{ISO})})$. For the cases of $n_{\text{iso}} = 0.963$ and $n_{\text{iso}} = 1$, we respectively obtain $(f_{\text{NL}}, \alpha^2 f_{\text{NL}}^{(\text{ISO})}) = (36 \pm 23, -39 \pm 69)$ and $(34 \pm 23, -22 \pm 70)$ without template marginalization. With template marginalization, these changes to $(f_{\text{NL}}, \alpha^2 f_{\text{NL}}^{(\text{ISO})}) = (32 \pm 23, 0 \pm 70)$ and $(38 \pm 24, -8 \pm 72)$. The error for each non-linearity parameter here is estimated by marginalizing over the other non-linearity parameter. In Fig. 1, we show 2D constraints in the $f_{\text{NL}} - \alpha^2 f_{\text{NL}}^{(\text{ISO})}$ plane for the cases with template marginalization. Due to the correlation of $\alpha^2 f_{\text{NL}}^{(\text{ISO})}$ with f_{NL} , a simultaneous fit for both of these variables changes the central value of $\alpha^2 f_{\text{NL}}^{(\text{ISO})}$, which are not so constrained as f_{NL} . We found that the dependence on n_{iso} is weak and the central values of $\alpha^2 f_{\text{NL}}^{(\text{ISO})}$ can change by 0.5σ by the treatment of the Galactic foregrounds.

Having all these results, we conclude that CMB data is consistent with Gaussianity at 2σ level, even if the uncorrelated CDM isocurvature perturbations are included in




Figure 1: Joint constraints on f_{NL} and $\alpha^2 f_{\text{NL}}^{(\text{ISO})}$ at 1 and 2σ levels with template marginalization. Solid red and dashed green lines respectively show the cases for $n_{\text{iso}} = 0.963$ and $n_{\text{iso}} = 1$, with the central values being indicated by the red plus and green cross. Blue star corresponds to the case where the perturbations are purely Gaussian.

consideration.

Let us compare our results with other studies. While our method is optimal based on the bispectrum, constraints on the same type of non-Gaussianity is studied in Ref. [48] based on the Minkowski functionals, which gives $\alpha^2 f_{\text{NL}}^{(\text{ISO})} = -15 \pm 60$, when the adiabatic perturbations are assumed to be Gaussian and $n_{\text{iso}} = 1$. We found our constraint is consistent with the previous study. On the other hand, we cannot find any improvement in the constraint, although our method is based on the optimal estimator and should be better than suboptimal ones. This suggests that the Minkowski functional method is almost optimal for uncorrelated CDM isocurvature perturbations. This can also be expected from the Fisher matrix analysis in Ref. [49], which gives a Cramér-Rao bound $\Delta(\alpha^2 f_{\text{NL}}) = 60$.

As a joint constraint on f_{NL} and $f_{\text{NL}}^{(\text{ISO})}$, our results are the first one obtained from observed data. On the other hand, the same constraint is forecasted using the Fisher matrix analysis in Ref. [49]. We note that 1σ errors of our results are consistent with the forecast.

5 Application to the axion model

In this section, we apply the constraints on the isocurvature non-Gaussianity to the axion isocurvature model. First, we shortly describe how non-Gaussian isocurvature perturbations in axion CDM arise in the inflationary Universe based on the δN -formalism [68, 69]. The baseline of our derivation is the same as in Ref. [32] (See also Ref. [48]). Throughout this section, we denote the energy density of a component i on the uniform density hyper-surface of the total matter with $\rho_i(\vec{x})$. We adopt the non-linear definition of isocurvature perturbations in a component i ,

$$S_i(\vec{x}) = 3(\zeta_i - \zeta)(\vec{x}), \quad (20)$$

where ζ_i and ζ are respectively the curvature perturbations on the uniform density hyper-surfaces of the component i and the total matter. According to the δN -formalism, $\rho(\vec{x})$ should be given by $\rho_i(\vec{x}) = \bar{\rho}_i e^{(1+w_i)S_i(\vec{x})}$, where $\bar{\rho}_i$ is the mean of $\rho_i(\vec{x})$ and w_i is the equation of state of the component i .

The axion is a pseudo Nambu-Goldstone boson of the Peccei-Quinn (PQ) U(1) symmetry, which solves the strong CP problem in quantum chromodynamics (QCD). If the PQ symmetry is broken during inflation, the axion has a classical field value $a_i = F_a \theta$ and a vacuum fluctuation δa , where F_a is the axion decay constant and $\theta \in [-\pi, \pi]$ is the initial misalignment angle during inflation. At high temperature, the axion is massless. As temperature T decreases the QCD phase transition takes place. At this moment, the axion becomes massive and starts the coherent oscillation around the true vacuum. This oscillation of the axion contributes to the energy density of CDM. We assume that CDM in the Universe is a mixture of the axion and other CDM components which are adiabatic.

The coherent oscillation of the axion synchronously starts on the uniform energy density hyper-surface of the total matter at around $m_{\text{axion}}(T) \simeq 3H(T)$ ^{#5}, where m_{axion} and H is the mass of axion and Hubble expansion rate, respectively. The energy density of the coherent oscillation is proportional to the square of its initial amplitude, $\rho_{\text{axion}}(\vec{x}) \propto (a_i + \delta a(\vec{x}))^2$, which leads

$$e^{S_{\text{axion}}(\vec{x})} = 1 + 2 \frac{a_i \delta a(\vec{x})}{a_*^2} + \frac{\delta a^2(\vec{x}) - \langle \delta a^2 \rangle}{a_*^2}, \quad (21)$$

where

$$a_*^2 \equiv a_i^2 + \langle \delta a^2 \rangle. \quad (22)$$

Since other CDM components are assumed to be adiabatic, their energy density is uniform on the uniform density hyper-surface of the total matter. Therefore $\rho_{\text{CDM}}(\vec{x})$ should be given by

$$\rho_{\text{CDM}}(\vec{x}) = \bar{\rho}_{\text{CDM}} \left[(1-r) + r e^{S_{\text{axion}}} \right], \quad (23)$$

^{#5}Here we assume that there are no isocurvature perturbations in neutrinos or, if any, extra radiations [3]. Otherwise, the coherent oscillation does not start synchronously on the uniform density hyper-surface and as a consequence, additional isocurvature perturbations in the axion can be induced.

where $r = \bar{\rho}_{\text{axion}}/\bar{\rho}_{\text{CDM}}$ is the energy fraction of the axion in CDM and $(1 - r)$ is that of other CDM components. Thus we finally obtain the isocurvature perturbations in CDM,

$$S_{\text{CDM}} = \ln \left[\frac{\rho_{\text{CDM}}(\vec{x})}{\bar{\rho}_{\text{CDM}}} \right] = \ln [1 - r + r e^{S_{\text{axion}}}]$$

$$\simeq 2r \frac{a_i \delta a}{a_*^2} + \left[\frac{1}{4r} \frac{a_*^2}{a_i^2} \right] \left[\left(2r \frac{a_i \delta a}{a_*^2} \right)^2 - \left\langle \left(2r \frac{a_i \delta a}{a_*^2} \right)^2 \right\rangle \right] + \dots \quad (24)$$

The last equality is approximately valid for $r \ll 1$. In the following, we keep the terms up to the second order.

Adopting the value of CDM abundance $\Omega_{\text{CDM}} h^2 = 0.11$ estimated by current observations [1], r is given by [70, 32]

$$r = 1.8 \left(\frac{F_a}{10^{12} \text{GeV}} \right)^{-0.82} \left[\left(\frac{F_a}{10^{12} \text{GeV}} \right)^2 \theta^2 + \left(\frac{H_{\text{inf}}/2\pi}{10^{12} \text{GeV}} \right)^2 \right], \quad (25)$$

where the first and second terms in the square bracket correspond to contributions from the classical field a_i and the fluctuation δa , respectively. The fluctuation of axion δa is almost scale-invariant (See also Ref. [12]) and its root mean square is given by $\sqrt{\langle \delta a^2 \rangle} = H_{\text{inf}}/2\pi$, where H_{inf} is the Hubble scale during inflation. Then power spectrum and bispectrum of CDM isocurvature perturbations are given by^{#6}

$$\langle S_{\text{CDM}}(\vec{k}_1) S_{\text{CDM}}(\vec{k}_2) \rangle = 4r^2 \frac{(H_{\text{inf}}/2\pi)^2}{(F_a \theta)^2 + (H_{\text{inf}}/2\pi)^2} \frac{2\pi^2}{k_1^3} (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2), \quad (26)$$

$$\langle S_{\text{CDM}}(\vec{k}_1) S_{\text{CDM}}(\vec{k}_2) S_{\text{CDM}}(\vec{k}_3) \rangle = 8r^3 \left[\frac{(H_{\text{inf}}/2\pi)^2}{(F_a \theta)^2 + (H_{\text{inf}}/2\pi)^2} \right]^2 \quad (27)$$

$$\times \left[\frac{(2\pi^2)^2}{k_1^3 k_2^3} + (2 \text{ perms}) \right] (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3).$$

Having all these results, constraints should be

$$1.8 \left(\frac{F_a}{10^{12} \text{GeV}} \right)^{-0.82} \left[\left(\frac{F_a}{10^{12} \text{GeV}} \right)^2 \theta^2 + \left(\frac{H_{\text{inf}}/2\pi}{10^{12} \text{GeV}} \right)^2 \right] < 1,$$

$$13 \left(\frac{F_a}{10^{12} \text{GeV}} \right)^{-1.64} \left[\left(\frac{F_a}{10^{12} \text{GeV}} \right)^2 \theta^2 + \left(\frac{H_{\text{inf}}/2\pi}{10^{12} \text{GeV}} \right)^2 \right] \left(\frac{H_{\text{inf}}/2\pi}{10^{12} \text{GeV}} \right)^2 < 0.15 A_s, \quad (28)$$

$$47 \left(\frac{F_a}{10^{12} \text{GeV}} \right)^{-2.46} \left[\left(\frac{F_a}{10^{12} \text{GeV}} \right)^2 \theta^2 + \left(\frac{H_{\text{inf}}/2\pi}{10^{12} \text{GeV}} \right)^2 \right] \left(\frac{H_{\text{inf}}/2\pi}{F_a \theta} \right)^4 < 140 A_s^2,$$

^{#6} Although, in the analysis in the previous sections, we omitted the loop contributions, we include them here. We also assume that logarithmic factors appear in the expression are considered to be $\mathcal{O}(1)$ and we here set them to be unity.

where $A_s = 2.43 \times 10^{-9}$ [1] is the amplitude of the adiabatic perturbations. From top to bottom, these three equations correspond to the constraints on the abundance $r < 1$, the isocurvature power spectrum $\alpha < 0.15$ (95 % CL) [1], and non-Gaussianity $|\alpha^2 f_{\text{NL}}^{(\text{ISO})}| < 140$ (2σ level) from our results. We should, however, note that since loop contributions are omitted in deriving the constraints in the previous sections, thus our results cannot in a rigorous manner be applied to cases for $F_a \theta \lesssim H_{\text{inf}}/2\pi$, where loop contributions dominate polyspectra. Therefore, in the parameter regions with $F_a \theta \lesssim H_{\text{inf}}/2\pi$, our constraints should be deemed as rough estimate.

In Fig. 2, we plotted the constraints of Eq. (28) in the $H_{\text{inf}}-\theta$ plane with several fixed values of F_a . Constraints have critical points at $F_a \theta = H_{\text{inf}}/2\pi$, where contributions of the classical field value and the fluctuation are comparable. We found that the constraint on non-Gaussianity from our results gives an upper bound on H_{inf} comparable to that from one on the isocurvature power spectrum.

When $H_{\text{inf}}/2\pi > F_a$, the PQ symmetry restores during the inflation^{#7} and in that case CDM axions are produced from the system of axionic string-wall system as well as the coherent oscillation with an initial alignment angle $\theta = \mathcal{O}(1)$. This excludes $H_{\text{inf}} \gtrsim 10^{11}\text{GeV}$ for $H_{\text{inf}}/2\pi > F_a$ in Fig. 2 (For details, we refer to e.g. Refs. [72, 73]).

These constraints lead that even if θ can be small by chance, large $F_a \gtrsim 10^{11}\text{GeV}$ is not allowed unless the inflation scale is low $H_{\text{inf}} \lesssim 10^{11}\text{GeV}$.

6 Conclusion

We studied constraints on non-Gaussianity in a mixture of adiabatic and uncorrelated CDM isocurvature perturbations. We adopted the optimal bispectrum estimator for the non-linearity parameters, and a fast method for non-Gaussian CMB simulation and the optimal weighting of observed maps are integrated in our analysis. Using the WMAP 7 year data of CMB temperature maps with template marginalization of the Galactic foregrounds, we obtained a constraint $\alpha^2 f_{\text{NL}}^{(\text{ISO})} = 40 \pm 62$ at 1σ level for the scale invariant isocurvature power spectrum when the adiabatic perturbations are assumed to be Gaussian. Under the same setup, we also obtained a joint constraint on the non-linearity parameters $(f_{\text{NL}}, \alpha^2 f_{\text{NL}}^{(\text{ISO})}) = (38 \pm 24, -8 \pm 72)$. The constraints weakly depend on the fiducial value of the isocurvature spectral index. Effects of the Galactic foregrounds at large angular scales are not severe. We found no statistically significant deviation from Gaussianity and the current WMAP observation of CMB is consistent with Gaussianity even when we include this type of isocurvature perturbations. We applied our results to the axion model.

Since the CDM and baryon isocurvature modes affect the CMB anisotropy in the same way except for the overall amplitude, our constraint can be translated into the baryon isocurvature perturbations. This can be easily done by substituting $\left(\frac{\Omega_b}{\Omega_c}\right)^3 \alpha^2 f_{\text{NL}}^{(\text{ISO})}$ for

^{#7}Here we assumed the reheating temperature T_{reh} is below H_{inf} . If $T_{\text{reh}} > H_{\text{inf}}$, the PQ symmetry can restore for smaller H_{inf} after inflation.




Figure 2: Constraints on axion isocurvature model in the $H_{\text{inf}}-\theta$ plane for several fixed values of F_a . Shaded regions are excluded by cosmological considerations. At small θ , the constraint on H_{inf} from the non-Gaussianity is slightly better than one from the power spectrum. See text for details.

$\alpha^2 f_{\text{NL}}^{(\text{ISO})}$, where Ω_b and Ω_c are the density parameters of baryon and CDM, respectively. Although we restrict ourselves to uncorrelated CDM isocurvature perturbations in this paper, our method can be generalized to the cases of non-Gaussianity in isocurvature perturbations correlated with adiabatic ones and other types such as neutrino ones. These will be studied in a forthcoming paper [52].

Acknowledgment

The authors would like to thank Eiichiro Komatsu and Kazunori Nakayama for helpful discussion. T. S. would like to thank the Japan Society for the Promotion of Science for the financial report. The authors acknowledge Kobayashi-Maskawa Institute for the Origin of Particles and the Universe, Nagoya University for providing computing resources in conducting the research reported in this paper. This work is supported by Grant-in-Aid for Scientific research from the Ministry of Education, Science, Sports, and Culture (MEXT), Japan, No. 14102004 (M.K.), No. 21111006 (M.K.), No. 23740195 (T.T.), No. 24740160 (C.H.) and also by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan. Some of the results in this paper have been derived using the HEALPix [67] package.

A Equivalence between $\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1}$ and $\langle S_{\text{prim}}^A S_{\text{prim}}^B \rangle_0$

In this appendix, we show that the normalization factor $\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1}$ and the covariance $\langle S_{\text{prim}}^A S_{\text{prim}}^B \rangle_0$ is equivalent.

From Eq. (15), we obtain

$$\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1} = \frac{1}{6} \sum_{\{lm\}} \hat{b}_{l_1 l_2 l_3}^A \mathcal{G}_{m_1 m_2 m_3}^{l_1 l_2 l_3} \langle \tilde{a}_{l_1 m_1} \tilde{a}_{l_2 m_2} \tilde{a}_{l_3 m_3} \rangle_{f_{\text{NL}}^B=1}, \quad (29)$$

where we used the fact that $\langle \tilde{a}_{lm} \rangle_{f_{\text{NL}}^A=1} = 0$. As discussed in Sec 3.3, we take $\tilde{a}_{lm} = \sum_{l'm'} [C^{-1}]_{lm, l'm'} a_{l'm'}$, so that

$$\begin{aligned} \langle \tilde{a}_{l_1 m_1} \tilde{a}_{l_2 m_2} \tilde{a}_{l_3 m_3} \rangle_{f_{\text{NL}}^B=1} &= \sum_{\{l'm'\}} [C^{-1}]_{l_1 m_1, l'_1 m'_1} [C^{-1}]_{l_2 m_2, l'_2 m'_2} [C^{-1}]_{l_3 m_3, l'_3 m'_3} \langle a_{l'_1 m'_1} a_{l'_2 m'_2} a_{l'_3 m'_3} \rangle_{f_{\text{NL}}^B=1} \\ &= \sum_{\{l'm'\}} [C^{-1}]_{l_1 m_1, l'_1 m'_1} [C^{-1}]_{l_2 m_2, l'_2 m'_2} [C^{-1}]_{l_3 m_3, l'_3 m'_3} \mathcal{G}_{m'_1 m'_2 m'_3}^{l'_1 l'_2 l'_3} \hat{b}_{l'_1 l'_2 l'_3}^B. \end{aligned} \quad (30)$$

Combining Eqs. (29) and (31), we obtain

$$\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1} = \frac{1}{6} \sum_{\{lm'l'm'\}} \hat{b}_{l_1 l_2 l_3}^A \mathcal{G}_{m_1 m_2 m_3}^{l_1 l_2 l_3} [C^{-1}]_{l_1 m_1, l'_1 m'_1} [C^{-1}]_{l_2 m_2, l'_2 m'_2} [C^{-1}]_{l_3 m_3, l'_3 m'_3} \mathcal{G}_{m'_1 m'_2 m'_3}^{l'_1 l'_2 l'_3} \hat{b}_{l'_1 l'_2 l'_3}^B. \quad (31)$$

On the other hand,

$$\begin{aligned} \langle S_{\text{prim}}^A S_{\text{prim}}^B \rangle_0 &= \frac{1}{36} \sum_{\{lm'l'm'\}} \hat{b}_{l_1 l_2 l_3}^A \hat{b}_{l'_1 l'_2 l'_3}^B \mathcal{G}_{m_1 m_2 m_3}^{l_1 l_2 l_3} \mathcal{G}_{m'_1 m'_2 m'_3}^{l'_1 l'_2 l'_3} \left[\langle \tilde{a}_{l_1 m_1} \tilde{a}_{l_2 m_2} \tilde{a}_{l_3 m_3} \tilde{a}_{l'_1 m'_1}^* \tilde{a}_{l'_2 m'_2}^* \tilde{a}_{l'_3 m'_3}^* \rangle_0 \right. \\ &\quad \left. - 6 \langle \tilde{a}_{l_1 m_1} \tilde{a}_{l'_1 m'_1}^* \tilde{a}_{l'_2 m'_2}^* \tilde{a}_{l'_3 m'_3}^* \rangle_0 \langle \tilde{a}_{l_2 m_2} \tilde{a}_{l_3 m_3} \rangle_0 + 9 \langle \tilde{a}_{l_1 m_1} \tilde{a}_{l'_1 m'_1}^* \rangle_0 \langle \tilde{a}_{l_2 m_2} \tilde{a}_{l_3 m_3} \rangle_0 \langle \tilde{a}_{l'_2 m'_2}^* \tilde{a}_{l'_3 m'_3}^* \rangle_0 \right]. \end{aligned} \quad (32)$$

By taking the Wick's contraction, the terms in the square bracket are reduced into

$$[C^{-1}]_{l_1 m_1, l'_1 m'_1} [C^{-1}]_{l_2 m_2, l'_2 m'_2} [C^{-1}]_{l_3 m_3, l'_3 m'_3} + (5 \text{ perms}), \quad (33)$$

where we used the fact that $\langle a_{lm} a_{l'm'} \rangle_0 = C_{lm, l'm'}$. Thus, we obtain

$$\langle S_{\text{prim}}^A S_{\text{prim}}^B \rangle_0 = \frac{1}{6} \sum_{\{lm'l'm'\}} \hat{b}_{l_1 l_2 l_3}^A \hat{b}_{l'_1 l'_2 l'_3}^B \mathcal{G}_{m_1 m_2 m_3}^{l_1 l_2 l_3} \mathcal{G}_{m'_1 m'_2 m'_3}^{l'_1 l'_2 l'_3} [C^{-1}]_{l_1 m_1, l'_1 m'_1} [C^{-1}]_{l_2 m_2, l'_2 m'_2} [C^{-1}]_{l_3 m_3, l'_3 m'_3}. \quad (34)$$

From Eqs. (31) and (34), we can finally see that $\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1} = \langle S_{\text{prim}}^A S_{\text{prim}}^B \rangle_0$.

B Estimation of normalization

Here we show how we can estimate the normalization factor $\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1}$ from simulation.

First we divide the CMB anisotropy a_{lm}^A into Gaussian and non-Gaussian parts

$$a_{lm}^A = a_{G,lm}^A + f_{\text{NL}}^A a_{\text{NG},lm}^A, \quad (35)$$

where $a_{G,lm}^A$ and $a_{\text{NG},lm}^A$ respectively are the Gaussian and non-Gaussian contributions, which are given by

$$a_{G,lm}^A = \int dr r^2 \alpha_l^A(r) \int d\hat{r} Y_{lm}^*(\hat{r}) X_G^A(\vec{r}), \quad (36)$$

$$a_{\text{NG},lm}^A = \int dr r^2 \alpha_l^A(r) \int d\hat{r} Y_{lm}^*(\hat{r}) (X_G^A(\vec{r})^2 - \langle X_G^A(\vec{r})^2 \rangle). \quad (37)$$

Since $\tilde{a}_{lm}$ is a linear combination of a_{lm} , we can also divide the weighted map into its Gaussian and non-Gaussian parts, $\tilde{a}_{lm}^A = \tilde{a}_{G,lm}^A + f_{\text{NL}}^A \tilde{a}_{\text{NG},lm}^A$. Then $\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1}$ in Eq. (29) can be rewritten as

$$\begin{aligned} \langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1} &= \frac{1}{6} \sum_{\{lm\}} \hat{b}_{l_1 l_2 l_3}^A \mathcal{G}_{m_1 m_2 m_3}^{l_1 l_2 l_3} \langle (\tilde{a}_{G,l_1 m_1} + \tilde{a}_{\text{NG},l_1 m_1}^B) (\tilde{a}_{G,l_2 m_2} + \tilde{a}_{\text{NG},l_2 m_2}^B) (\tilde{a}_{G,l_3 m_3} + \tilde{a}_{\text{NG},l_3 m_3}^B) \rangle_{f_{\text{NL}}^B=1} \\ &= \frac{1}{6} \sum_{\{lm\}} \hat{b}_{l_1 l_2 l_3}^A \mathcal{G}_{m_1 m_2 m_3}^{l_1 l_2 l_3} (3 \langle \tilde{a}_{\text{NG},l_1 m_1}^B \tilde{a}_{G,l_2 m_2}^B \tilde{a}_{G,l_3 m_3}^B \rangle + \langle \tilde{a}_{\text{NG},l_2 m_2}^B \tilde{a}_{\text{NG},l_2 m_2}^B \tilde{a}_{\text{NG},l_3 m_3}^B \rangle)_{f_{\text{NL}}^B=1}, \end{aligned} \quad (38)$$

where the last equality follows from the Wick's theorem. As long as non-Gaussianity is not too large, the second term in the last line can be omitted. Although we need many MC samples to evaluate the term $\langle \tilde{a}_{\text{NG},l_1m_1}^B \tilde{a}_{G,l_2m_2}^B \tilde{a}_{G,l_3m_3}^B \rangle$, we checked that $\mathcal{O}(10)$ MC samples are enough for this purpose. In addition, since the terms in the last line of Eq. (38) consists of only anisotropy $\tilde{a}_{lm}^B$ from a single perturbation mode X^B , we can estimate $\langle S_{\text{prim}}^A \rangle_{f_{\text{NL}}^B=1}$ without assuming a fiducial value for the fraction of the isocurvature contribution α .

References

- [1] E. Komatsu *et al.* [WMAP Collaboration], *Astrophys. J. Suppl.* **192**, 18 (2011) [arXiv:1001.4538 [astro-ph.CO]].
- [2] M. Bucher, K. Moodley and N. Turok, *Phys. Rev. D* **62**, 083508 (2000) [astro-ph/9904231].
- [3] M. Kawasaki, K. Miyamoto, K. Nakayama and T. Sekiguchi, *JCAP* **1202**, 022 (2012) [arXiv:1107.4962 [astro-ph.CO]].
- [4] E. Kawakami, M. Kawasaki, K. Miyamoto, K. Nakayama and T. Sekiguchi, *JCAP* **1207**, 037 (2012) [arXiv:1202.4890 [astro-ph.CO]].
- [5] M. Axenides, R. H. Brandenberger, M. S. Turner, *Phys. Lett.* **B126**, 178 (1983).
- [6] D. Seckel, M. S. Turner, *Phys. Rev.* **D32**, 3178 (1985).
- [7] A. D. Linde, *Phys. Lett.* **B158**, 375-380 (1985).
- [8] A. D. Linde, D. H. Lyth, *Phys. Lett.* **B246**, 353-358 (1990).
- [9] M. S. Turner, F. Wilczek, *Phys. Rev. Lett.* **66**, 5 (1991).
- [10] A. D. Linde, *Phys. Lett.* **B259**, 38 (1991).
- [11] D. H. Lyth, *Phys. Rev.* **D45**, 3394 (1992).
- [12] S. Kasuya, M. Kawasaki, *Phys. Rev.* **D80**, 023516 (2009) [arXiv:0904.3800 [astro-ph.CO]].
- [13] K. Enqvist and J. McDonald, *Phys. Rev. Lett.* **83**, 2510 (1999) [arXiv:hep-ph/9811412].
- [14] K. Enqvist, J. McDonald, *Phys. Rev.* **D62**, 043502 (2000) [hep-ph/9912478].
- [15] M. Kawasaki and F. Takahashi, *Phys. Lett. B* **516**, 388 (2001) [arXiv:hep-ph/0105134].

- [16] K. Enqvist and M. S. Sloth, Nucl. Phys. B **626**, 395 (2002) [arXiv:hep-ph/0109214].
- [17] D. H. Lyth and D. Wands, Phys. Lett. B **524**, 5 (2002) [arXiv:hep-ph/0110002].
- [18] T. Moroi and T. Takahashi, Phys. Lett. B **522**, 215 (2001) [Erratum-ibid. B **539**, 303 (2002)] [arXiv:hep-ph/0110096].
- [19] T. Moroi and T. Takahashi, Phys. Rev. D **66**, 063501 (2002) [arXiv:hep-ph/0206026].
- [20] D. H. Lyth, C. Ungarelli and D. Wands, Phys. Rev. D **67**, 023503 (2003) [arXiv:astro-ph/0208055].
- [21] T. Moroi and H. Murayama, Phys. Lett. B **553**, 126 (2003) [hep-ph/0211019].
- [22] D. H. Lyth and D. Wands, Phys. Rev. D **68**, 103516 (2003) [arXiv:astro-ph/0306500].
- [23] K. Hamaguchi, M. Kawasaki, T. Moroi and F. Takahashi, Phys. Rev. D **69**, 063504 (2004) [hep-ph/0308174].
- [24] C. Gordon and K. A. Malik, Phys. Rev. D **69**, 063508 (2004) [astro-ph/0311102].
- [25] M. Ikegami and T. Moroi, Phys. Rev. D **70**, 083515 (2004) [hep-ph/0404253].
- [26] F. Ferrer, S. Rasanen and J. Valiviita, JCAP **0410**, 010 (2004) [astro-ph/0407300].
- [27] M. Beltran, Phys. Rev. D **78**, 023530 (2008) [arXiv:0804.1097 [astro-ph]].
- [28] T. Moroi and T. Takahashi, Phys. Lett. B **671**, 339 (2009) [arXiv:0810.0189 [hep-ph]].
- [29] M. Lemoine, J. Martin and J. Yokoyama, Europhys. Lett. **89**, 29001 (2010) [arXiv:0903.5428 [astro-ph.CO]].
- [30] M. Lemoine, J. Martin and J. Yokoyama, Phys. Rev. D **80**, 123514 (2009) [arXiv:0904.0126 [astro-ph.CO]].
- [31] T. Takahashi, M. Yamaguchi and S. Yokoyama, Phys. Rev. D **80**, 063524 (2009) [arXiv:0907.3052 [astro-ph.CO]].
- [32] M. Kawasaki, K. Nakayama, T. Sekiguchi, T. Suyama and F. Takahashi, JCAP **0811**, 019 (2008) [arXiv:0808.0009 [astro-ph]].
- [33] D. Langlois, F. Vernizzi and D. Wands, JCAP **0812**, 004 (2008) [arXiv:0809.4646 [astro-ph]].
- [34] M. Kawasaki, K. Nakayama, T. Sekiguchi, T. Suyama and F. Takahashi, JCAP **0901**, 042 (2009) [arXiv:0810.0208 [astro-ph]].
- [35] E. Kawakami, M. Kawasaki, K. Nakayama and F. Takahashi, JCAP **0909**, 002 (2009) [arXiv:0905.1552 [astro-ph.CO]].

- [36] D. Langlois and A. Lepidi, JCAP **1101**, 008 (2011) [arXiv:1007.5498 [astro-ph.CO]].
- [37] D. Langlois and T. Takahashi, JCAP **1102**, 020 (2011) [arXiv:1012.4885 [astro-ph.CO]].
- [38] K. Moodley, M. Bucher, J. Dunkley, P. G. Ferreira and C. Skordis, Phys. Rev. D **70**, 103520 (2004) [astro-ph/0407304].
- [39] M. Beltran, J. Garcia-Bellido, J. Lesgourgues and A. Riazuelo, Phys. Rev. D **70**, 103530 (2004) [astro-ph/0409326].
- [40] R. Bean, J. Dunkley and E. Pierpaoli, Phys. Rev. D **74**, 063503 (2006) [astro-ph/0606685].
- [41] R. Trotta, Mon. Not. Roy. Astron. Soc. **375**, L26 (2007) [astro-ph/0608116].
- [42] R. Kesitalo, H. Kurki-Suonio, V. Muhonen and J. Valiviita, JCAP **0709**, 008 (2007) [astro-ph/0611917].
- [43] M. Kawasaki and T. Sekiguchi, Prog. Theor. Phys. **120**, 995 (2008) [arXiv:0705.2853 [astro-ph]].
- [44] I. Sollom, A. Challinor and M. P. Hobson, Phys. Rev. D **79**, 123521 (2009) [arXiv:0903.5257 [astro-ph.CO]].
- [45] J. Valiviita and T. Giannantonio, Phys. Rev. D **80**, 123516 (2009) [arXiv:0909.5190 [astro-ph.CO]].
- [46] H. Li, J. Liu, J. -Q. Xia and Y. -F. Cai, Phys. Rev. D **83**, 123517 (2011) [arXiv:1012.2511 [astro-ph.CO]].
- [47] J. Valiviita, M. Savelainen, M. Talvitie, H. Kurki-Suonio and S. Rusak, Astrophys. J. **753**, 151 (2012) [arXiv:1202.2852 [astro-ph.CO]].
- [48] C. Hikage, K. Koyama, T. Matsubara, T. Takahashi and M. Yamaguchi, Mon. Not. Roy. Astron. Soc. **398**, 2188 (2009) [arXiv:0812.3500 [astro-ph]].
- [49] C. Hikage, D. Munshi, A. Heavens and P. Coles, Mon. Not. Roy. Astron. Soc. **404**, 1505 (2010) [arXiv:0907.0261 [astro-ph.CO]].
- [50] D. Langlois and B. van Tent, Class. Quant. Grav. **28**, 222001 (2011) [arXiv:1104.2567 [astro-ph.CO]].
- [51] D. Langlois and B. van Tent, JCAP **1207**, 040 (2012) [arXiv:1204.5042 [astro-ph.CO]].
- [52] C. Hikage, M. Kawasaki, T. Sekiguchi and T. Takahashi, in preparation.
- [53] E. Komatsu and D. N. Spergel, Phys. Rev. D **63**, 063002 (2001) [astro-ph/0005036].

- [54] K. M. Smith, L. Senatore and M. Zaldarriaga, JCAP **0909**, 006 (2009) [arXiv:0901.2572 [astro-ph]].
- [55] E. Komatsu, D. N. Spergel and B. D. Wandelt, Astrophys. J. **634**, 14 (2005) [astro-ph/0305189].
- [56] A. P. S. Yadav, E. Komatsu and B. D. Wandelt, Astrophys. J. **664**, 680 (2007) [astro-ph/0701921].
- [57] P. Creminelli, A. Nicolis, L. Senatore, M. Tegmark and M. Zaldarriaga, JCAP **0605**, 004 (2006) [astro-ph/0509029].
- [58] A. P. S. Yadav, E. Komatsu, B. D. Wandelt, M. Liguori, F. K. Hansen and S. Matarrese, Astrophys. J. **678**, 578 (2008) [arXiv:0711.4933 [astro-ph]].
- [59] M. Liguori, S. Matarrese and L. Moscardini, Astrophys. J. **597**, 57 (2003) [astro-ph/0306248].
- [60] M. Liguori, A. Yadav, F. K. Hansen, E. Komatsu, S. Matarrese and B. Wandelt, Phys. Rev. D **76**, 105016 (2007) [Erratum-ibid. D **77**, 029902 (2008)] [arXiv:0708.3786 [astro-ph]].
- [61] F. Elsner and B. D. Wandelt, Astrophys. J. Suppl. **184**, 264 (2009) [arXiv:0909.0009 [astro-ph.CO]].
- [62] S. P. Oh, D. N. Spergel and G. Hinshaw, Astrophys. J. **510**, 551 (1999) [astro-ph/9805339].
- [63] K. M. Smith, O. Zahn and O. Dore, Phys. Rev. D **76**, 043510 (2007) [arXiv:0705.3980 [astro-ph]].
- [64] A. Lewis, A. Challinor and A. Lasenby, Astrophys. J. **538**, 473 (2000) [astro-ph/9911177].
- [65] N. Jarosik, C. L. Bennett, J. Dunkley, B. Gold, M. R. Greason, M. Halpern, R. S. Hill and G. Hinshaw *et al.*, Astrophys. J. Suppl. **192**, 14 (2011) [arXiv:1001.4744 [astro-ph.CO]].
- [66] B. Gold, N. Odegard, J. L. Weiland, R. S. Hill, A. Kogut, C. L. Bennett, G. Hinshaw and J. Dunkley *et al.*, Astrophys. J. Suppl. **192**, 15 (2011) [arXiv:1001.4555 [astro-ph.GA]].
- [67] K. M. Gorski, E. Hivon, A. J. Banday, B. D. Wandelt, F. K. Hansen, M. Reinecke and M. Bartelman, Astrophys. J. **622**, 759 (2005) [astro-ph/0409513].
- [68] M. Sasaki and E. D. Stewart, Prog. Theor. Phys. **95**, 71 (1996) [astro-ph/9507001].

- [69] D. H. Lyth, K. A. Malik and M. Sasaki, JCAP **0505**, 004 (2005) [astro-ph/0411220].
- [70] M. S. Turner, Phys. Rev. D **33**, 889 (1986).
- [71] D. H. Lyth, JCAP **0712**, 016 (2007) [arXiv:0707.0361 [astro-ph]].
- [72] T. Hiramatsu, M. Kawasaki, T. Sekiguchi, M. Yamaguchi and J. 'i. Yokoyama, Phys. Rev. D **83**, 123531 (2011) [arXiv:1012.5502 [hep-ph]].
- [73] T. Hiramatsu, M. Kawasaki, K. 'i. Saikawa and T. Sekiguchi, Phys. Rev. D **85**, 105020 (2012) [arXiv:1202.5851 [hep-ph]].